

In The Claims:

1. (Currently Amended) ~~An A monoclonal antibody against a fusion polypeptide comprising a histidine portion, wherein said antibody is directed against binds specifically to said histidine portion but not to the non-histidine portion of the fusion polypeptide, and wherein said histidine portion comprises 6-18 successive histidine residues.~~
2. (Currently Amended) The monoclonal antibody of claim 1, wherein said antibody is a polyclonal antibody said histidine portion comprises 6 successive histidine residues.
3. (Canceled)
4. (Canceled)
5. (Withdrawn) A process for the preparation of the polyclonal antibody of claim 2, comprising:
 - (a) immunizing an animal with a histidine fusion polypeptide; and
 - (b) collecting said polyclonal antibody from the serum of said animal.
6. (Withdrawn) The process of claim 5, wherein a mixture of different histidine fusion polypeptides is used for immunization.
7. (Withdrawn) A method for detecting a fusion polypeptide having a histidine portion, comprising:
 - (a) incubating said polypeptide with the antibody of Claim 1, 2, 3, or 4; and
 - (b) detecting the antibody in a detection reaction.

Appln. No.: 10/686,355; Filed: October 14, 2003
Reply to Non-Final Office Action mailed July 7, 2006
Response dated November 6, 2006

8. (Withdrawn) The method of claim 7, wherein the detection reaction is selected from the group consisting of Western blot, ELISA, immunofluorescence, and immunoprecipitation.

9. (Withdrawn) A process for the preparation of the monoclonal antibody of claim 3, comprising;

- (a) immunizing an animal with a histidine fusion polypeptide;
- (b) fusing the animal's spleen cells with myeloma cells to generate hybridoma cells; and
- (c) obtaining said monoclonal antibody form said hybridoma cells.

10. (Withdrawn) The process of claim 9, wherein a mixture of different histidine fusion polypeptides is used for immunization.